



King's Research Portal

DOI:

[10.1111/add.13893](https://doi.org/10.1111/add.13893)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Raw, M., Ayo-Yusuf, O., Chaloupka, F., Fiore, M., Glynn, T., Hawari, F., Mackay, J., McNeill, A. D., & Reddy, S. (2017). Recommendations for the implementation of WHO Framework Convention on Tobacco Control Article 14 on tobacco cessation support. *Addiction*. <https://doi.org/10.1111/add.13893>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Recommendations for the implementation of WHO Framework Convention on Tobacco Control Article 14 on tobacco cessation support

Martin Raw, Visiting Professor, Department of Population Health, New York University School of Medicine, New York, USA and UK Centre for Tobacco and Alcohol Studies, Division of Epidemiology and Public Health, University of Nottingham, Nottingham, UK

Olalekan Ayo-Yusuf, Professor and Dean, Faculty of Health Sciences, Sefako Makgatho Health Sciences University, Pretoria, South Africa

lekan.ayo-yusuf@smu.ac.za

Frank Chaloupka, Director, Health Policy Center, Institute for Health Research and Policy, University of Illinois at Chicago, Chicago, USA

fjc@uic.edu

Michael Fiore, Director, Center for Tobacco Research and Intervention and Professor of Medicine, University of Wisconsin School of Medical and Public Health, Madison, USA

mcf@ctri.medicine.wisc.edu

Thomas Glynn, Consulting Professor, Stanford Prevention Research Center, School of Medicine, Stanford University, Palo Alto, USA

tomglynn7@gmail.com

Feras Hawari, Director, Cancer Control Office and Chief, Section of Pulmonary and Critical Care, King Hussein Cancer Center, Amman, Jordan

fhawari@khcc.jo

Judith Mackay, Senior Advisor, Vital Strategies, and Director, Asian Consultancy on Tobacco Control, Hong Kong

jmackay1@netvigator.com

Ann McNeill, Professor of Tobacco Addiction, UK Centre for Tobacco & Alcohol Studies, National Addiction Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

ann.mcneill@kcl.ac.uk

Srinath Reddy, President, Public Health Foundation of India, Delhi NCR, Gurgaon, Haryana, India

ksrinath.reddy@phfi.org

Corresponding author: Martin Raw martin@martinraw.com

Word count: 2620

Key words: tobacco cessation, tobacco dependence treatment, FCTC Article 14

Article 14 of the Framework Convention for Tobacco Control **recommends** provision of evidence-based support for tobacco cessation. However, it is being implemented very slowly in most countries, so that by 2014 only a small minority of the world's tobacco users have access to appropriate cessation support, according to WHO. This editorial, endorsed by more than **60** global leaders in tobacco cessation (see online version), sets out actions that countries can take to remedy the situation, focusing on low cost, broad reach interventions.

Introduction

Stopping tobacco use is a vital element in any comprehensive approach to tobacco control. It saves lives, prevents debilitating illnesses and is beneficial economically (3), not least by reducing healthcare costs (4). Recognizing this, the World Health Organization oversaw the development of the world's first global health treaty, the Framework Convention on Tobacco Control (WHO FCTC), which entered into force in 2005, and includes an Article on tobacco cessation (Article 14). The FCTC is one of the most widely supported of all United Nations (UN) treaties. At the time of writing 179 countries (180 Parties), representing 90% of the world's population, have ratified the Treaty (1). In 2010 the fourth Conference of the Parties to the Treaty adopted guidelines to assist Parties in implementing Article 14 of the Treaty, on tobacco dependence and cessation (2).

Since 2010, several important international initiatives have provided further impetus for the implementation of the FCTC and Article 14 of the Treaty:

- In 2011 the United Nations General Assembly held a special session at which it adopted a political declaration on non-communicable diseases (NCDs), which recognized tobacco as one of the four main NCD risk factors and committed governments to accelerated implementation of the FCTC (5).
- In 2013 the World Health Assembly adopted voluntary global targets on NCDs, including a 25% decrease in premature mortality from NCDs by 2025 and a 30% relative reduction in tobacco use by the same year (6) compared with a 2010 baseline.
- In 2015 world leaders agreed to an ambitious set of Sustainable Development Goals (SDGs). Goal 3 ("Ensure healthy lives and promote well-being for all at all ages") includes a target of reducing premature mortality from NCDs by 33% by 2030, and goal 3a is to strengthen implementation of the FCTC in all countries (7).

Thus there is strong international support to strengthen and accelerate full implementation of the WHO FCTC. While preventing people from starting smoking and using other forms of tobacco will save lives twenty years and more from now, only effective tobacco cessation will have a sufficient effect on mortality in the few years left to reach the UN/WHO goal of a 25% reduction in premature mortality from NCDs by 2025. Failure on that front will surely thwart the global aspiration for success in NCD control and fail to save millions of current smokers' lives (8).

A neglected Article

The 2014 WHO Global Progress Report notes that in general good progress is being made in FCTC implementation (although it notes that only 19% of Parties have implemented it to the "highest level") but that some Treaty articles are being implemented more quickly than others (9). The report shows that by 2014 the strongest focus was on Articles 8 (restrictions on smoking in public places), 16 (restrictions on sales to minors), 11 (large health warnings) and 12 (educational programmes), which motivate tobacco users to seek treatment for tobacco dependence. Only half of Parties reported that they were implementing Article 14, on tobacco dependence and cessation, a finding consistent with that of another survey suggesting slow implementation of Article 14 (10). Evidence from the survey

suggests that perceived cost may be one reason for hesitation in addressing tobacco treatment support. Another reason may be lack of clarity about the effectiveness and cost effectiveness of tobacco dependence treatment.

There is little doubt that implementation of FCTC Article 14 and its guidelines will bring health and economic benefits to countries (3). Helping smokers to stop is a highly cost effective healthcare intervention which saves lives (11, 12, 13, 14, 15), improves population health by reducing both morbidity and mortality (16, 17, 18) and can reduce health-care costs (4). Every day that smokers over 35 years old continue to smoke they lose about 3–6 hours of life (19). So for the world's estimated 1 billion current smokers (3), about half of them over 35 (20), 62 million days of life are lost *every day*. Some of these will stop unaided. Many others will stop only after repeated attempts over time. Many more smokers will never stop and will die before they can stop (21), mainly because tobacco use is so addictive (2, 22, 23, 24). Arguably treatment support should be provided along with measures like restrictions on smoking in public places and other measures mentioned in the FCTC (25). While other tobacco control policies, such as tax increases and smokefree air laws, primarily increase quit attempts, evidence shows that comprehensive cessation treatment policies primarily affect quit success (12). Combining cessation treatment with other policies creates a synergistic impact that improves the effect of other policies. The unaided tobacco cessation rate at six to twelve months is low, only about 3–5% (26). Adding comprehensive cessation support can increase this considerably, at least doubling it (12, 18, 27, 28), but even brief advice improves cessation rates significantly. Its long term rate is much lower, but applied at population level it would still result in significant population health gain (27). One reason why so many people who eventually end up as ex-smokers have stopped without help is not because attempting to quit without help is the most effective method, but simply because many of them had no access to help. If they had had access to support, they may have stopped sooner, and many life years could have been saved. Additionally, many smokers who never stopped might have managed to do so if support had been available and accessible. According to WHO just 15% of the world's population have access to appropriate cessation support (29), and even among this 15% many do not use the support that is available (30). And although less is known about smokeless tobacco use clearly a comprehensive approach is required for cessation (31, 32).

There is thus a significant unmet need for tobacco cessation support. Although some tobacco users do stop unaided, many more would stop if they had access to effective support. The FCTC Article 14 Guidelines recommend how to address this need in a way that is appropriate to the situation of the country, and which makes best use of existing resources. Improving cessation support need not be expensive, and helping smokers stop sooner rather than later will not only save many lives but improve the quality of life of those who do stop.

Because of these considerations, and because the FCTC Article 14 Guidelines contain clear and feasible recommendations, we believe the time has come to re-evaluate the role of cessation support, especially because effective, low-cost, broad-reach approaches exist that can be implemented quickly (27). Revenue could be raised for tobacco control and cessation from tobacco tax increases (3, 33, 34) and evidence from some countries suggests that public support for such tax increases is stronger if some of the money is spent offering support to tobacco users who need it, many of whom are poor and for whom tobacco use is a financial strain (35).

Some underlying principles

The FCTC Article 14 Guidelines include the following key underlying principles:

- Implementing cessation support measures in conjunction with population level interventions

- covered by other FCTC articles will have a synergistic effect and maximize their impact.
- Some Parties will want or need to follow a stepwise approach in developing tobacco cessation support that focuses first on what is feasible and affordable, and then on what can be added when more resources are available.
 - To develop tobacco dependence treatment as rapidly as possible and at as low a cost as possible, Parties should use existing resources, workforce and infrastructure, including for example community health workers, frontline primary care providers, and tuberculosis services.
 - Parties are encouraged to create a sustainable infrastructure which motivates attempts to quit, ensures wide access to support, and provides sustainable resources for such support, which should be affordable.
 - Tobacco cessation support is a key component of a *comprehensive*, integrated tobacco control programme; offering tobacco users support for their cessation efforts will reinforce other tobacco control policies, by increasing support for them and increasing their acceptability.
 - Healthcare systems should play a central role in the delivery of cessation support.
 - Parties should follow the evidence base and also keep up with new developments.
 - Tobacco cessation and treatment strategies should be based on the best available evidence; there is clear scientific evidence that tobacco dependence treatment is effective, is a cost effective healthcare intervention, and thus a worthwhile investment for healthcare systems.
 - Tobacco cessation and treatment strategies should be as inclusive as possible, considering the needs of vulnerable groups.

Recommendations

FCTC Article 14 states that “each Party shall develop and disseminate appropriate, comprehensive and integrated guidelines based on scientific evidence and best practices, taking into account national circumstances and priorities, and shall take effective measures to promote cessation of tobacco use and adequate treatment for tobacco dependence”.

The FCTC Article 14 Guidelines set out detailed recommendations for implementation of this Article, including on developing infrastructure, the key components of a national treatment system, and a stepwise approach to developing cessation support. We urge all stakeholders to use these guidelines to inform their thinking about cessation support.

Eventually all Parties should implement all aspects of the FCTC, and the FCTC Article 14 Guidelines encourage Parties to implement measures beyond those recommended by the guidelines (in accordance with the provisions of Article 2.1 of the Convention). However they also clearly acknowledge that different countries will be at different stages and will need to implement Article 14 according to their own priorities and resources. Thus we recommend measures we believe all countries can begin implementing now, then measures to be implemented as resources become available.

We recommend that countries consider implementing *these six core measures* immediately:

- Conduct a national situation analysis, as recommended in the FCTC Article 14 Guidelines.
- Develop and implement an evidence based national cessation strategy and national cessation guidelines.
- Mandate recording tobacco use of all patients in all medical notes.
- Train healthcare workers to record tobacco use and give brief advice.
- Help healthcare workers to quit tobacco use.

- Integrate brief advice to all tobacco users into the healthcare system.

Some of these measures are relatively low cost (27), and we believe some are essential if healthcare systems are to play an active role in helping tobacco users stop. For example, healthcare workers raising the issue and giving brief advice will be more likely if tobacco use is recorded in medical notes, yet only 20% of countries mandate this (10). The Article 14 Guidelines strongly emphasise using existing infrastructure as far as possible, both to keep costs down and to ensure as broad coverage as possible. They mention infrastructure like the primary healthcare system, and services to treat tuberculosis and HIV/AIDS, and in some countries dental services and NCD programmes could also be involved. The first WHO Report on the Global Tobacco Epidemic (the MPOWER package) recommends that tobacco cessation advice be incorporated into primary healthcare services (36). We believe that whatever stage countries are at in implementing other articles of the FCTC, they could be implementing these six core measures.

We recommend that as resources become available countries implement these additional measures:

- Offer mobile phone text messaging support and other evidence based web support (very low cost and accessible).
- Make evidence based affordable cessation medicines and less harmful forms of nicotine more available.
- Offer evidence based telephone quitlines and put the number on pack warnings.
- Expand training to include other healthcare workers, for example, village health workers.
- Offer specialist face-to-face cessation support.
- Establish mass communication and education programmes that encourage cessation and promote cessation support.

Keeping up-to-date with the scientific evidence

The FCTC Article 14 Guidelines urge Parties to follow the evidence base, keep under review the emerging scientific evidence, and be open to innovative approaches to promote tobacco cessation (2). Since the Article 14 Guidelines were adopted in 2010 for example, new research has emerged on text messaging and on less harmful forms of nicotine delivery. Thus national cessation guidelines need to be reviewed and updated periodically to include new interventions. An increasing body of evidence demonstrates the effectiveness of text messaging, a measure with enormous potential because of its potential population reach and low cost. Research is also developing on different forms of nicotine delivery. Clearly any cessation guideline published in 1980 would be seriously out-of-date by 1990 because of the advent of nicotine replacement therapy (NRT) – itself an alternative form of nicotine delivery when it was first licensed – now affordable in many countries, and on the WHO Essential Medicines List (37). Two recent authoritative publications recognise the continuum of risk from using nicotine products (38, 39). Based on this idea, differential taxes have been proposed that discourage use of the most dangerous forms of nicotine yielding products (40). We therefore urge countries to remember that it is the nicotine in tobacco products that keeps people using them, but it is primarily the toxins in the tobacco and tobacco smoke that cause the illness and death, and to keep under review all potential options for tobacco cessation, including, for some, the continued use of nicotine.

Focusing on affordability

Many tools exist to help countries implement or improve their provision of tobacco cessation support, including the recommendations of the FCTC Article 14 Guidelines themselves (for example to include the quitline number on tobacco packaging) (2), WHO resources (41, 42, 43),

and tools to help countries conduct a national situation analysis – recommended in the Article 14 Guidelines – and access other related resources (44). West and colleagues have published a review of effectiveness and affordability, which includes an excel calculator that permits a country to input its own national data in order to calculate the affordability of a cessation intervention (27). Potentially low cost evidence based medications are available (45), and countries should also examine how to make medications more affordable, through bulk buying for example, and by including some on their national essential medicines lists. Many countries base their national lists on WHO's essential medicines list (37), which includes nicotine chewing gum and nicotine patch. We also urge countries to consider increasing funding for tobacco control and cessation support through earmarked tax increases (34), a measure recommended in the FCTC Article 6 Guidelines.

Obviously a key challenge in providing specialist face-to-face cessation support is cost, but another issue is reach. Even in some countries that do provide specialist support it is only accessible to a small fraction of the population. This is why the Article 14 Guidelines encourage Parties to *consider using existing, rather than creating new, infrastructures*. However many countries do have the resources to offer specialist support, and the Article 14 guidelines include a number of recommendations for these, including that it could be delivered by a variety of healthcare or other trained workers, in a wide variety of settings, should be easily accessible to tobacco users, and where possible should be provided free or at an affordable cost. The guidelines also emphasize the importance of mass communication programmes that promote cessation.

Conclusions

Implementation of FCTC Article 14 has been slow. Accelerating its implementation to complement implementation of other Articles, including those promoting tax increases, smokefree public places and graphic health warnings, will save lives, prevent numerous debilitating illnesses, and bring economic benefits. The availability of effective, broad-reach, low-cost interventions, and methods to help countries select affordable treatments (27) will remove significant barriers to the development of tobacco cessation support and make Article 14 implementation both eminently feasible and a significant contributor to global health.

Acknowledgements

We gratefully acknowledge help from Brent Bell, Alice Grainger-Gasser, Ehsan Latif, Kapka Nilan, Kamran Siddiqi and Francis Thompson in the preparation of this paper.

Author contributions

MR led the writing of this article with the support and active involvement of all authors.

Author interests statements

MR was funded by Global Bridges, Mayo Clinic, to lead the writing of this article; TG was funded by Global Bridges, Mayo Clinic during the writing of this article; OAY, FC, MF, FH, JM, AM and SR declare no conflicts of interest. The Global Bridges project at Mayo Clinic is partially funded by Pfizer, but managed independently.

Terminology

The terms tobacco dependence treatment and tobacco cessation support are used interchangeably here and in the FCTC Article 14 Guidelines, which also note that treatment includes broad reach

low intensity population approaches, behavioural support, and effective medications. This article is about stopping smoking and other forms of tobacco use.

References

1. World Health Organization. WHO Framework Convention on Tobacco Control. Geneva: World Health Organization 2005.
2. FCTC Article 14 Guidelines <http://www.who.int/fctc/Guidelines.pdf> (accessed 20 Feb 17)
3. US National Cancer Institute and World Health Organization. *The Economics of Tobacco and Tobacco Control*. NCI Tobacco Control Monograph 21. NIH Publication No 16-CA-8029A. Bethesda, MD: US Department of Health and Human Services, National Institutes of Health, National Cancer Institute and Geneva: World Health Organization 2016.
4. National Institute for Health and Care Excellence. Tobacco return on investment tool. <https://www.nice.org.uk/about/what-we-do/into-practice/return-on-investment-tools/tobacco-return-on-investment-tool> (accessed 28 March 2017).
5. President of the General Assembly of the United Nations. Political declaration of the high-level meeting on the prevention and control of non-communicable diseases. Sixty sixth session, 16 September 2011, (accessed 28 March 2017) http://www.who.int/entity/nmh/events/un_ncd_summit2011/political_declaration_en.pdf
6. World Health Organization. Global Action Plan for the prevention and control of noncommunicable diseases 2013-2020. Geneva, World Health Organization, 2013
7. United Nations General Assembly, Sixty ninth session. Transforming our world: The 2030 agenda for Sustainable Development. 12 August 2015 (accessed 28 March 2017) <https://sustainabledevelopment.un.org/post2015/transformingourworld>
8. Kontis V, Mathers CD, Rehm J, et al. Contribution of six risk factors to achieving the 25×25 non-communicable disease mortality reduction target: a modelling study. *Lancet* 2014; 384: 427–37.
9. WHO FCTC Secretariat. 2014 Global progress report on implementation of the Framework Convention on Tobacco Control. Geneva, WHO Framework Convention on Tobacco Control Secretariat, 2014.
10. Piné-Abata H, McNeill A, Murray R, Bitton A, Rigotti N, Raw M. A survey of tobacco dependence treatment services in 121 countries. *Addiction* 2013; 108: 1476–84.
11. Levy DT, Mabry PL, Graham AL, Abrams DB, Graham AL, Orleans CT. Reaching Healthy People 2010 by 2013: A SimSmoke Simulation. *American Journal of Preventive Medicine* 2010;38(38):S373-81.
12. Levy DT, Graham AL, Mabry PL, Abrams DB, Graham AL, Orleans CT. Modeling the Impact of Smoking Cessation Treatment Policies on Quit Rates. *American Journal of Preventive Medicine* 2010;38(38):S364-72.
13. Levy DT, Friend K. Examining the effects of tobacco treatment policies on smoking rates and smoking related deaths using the SimSmoke computer simulation model. *Tobacco Control* 2002;11:47-54.
14. Levy DT, Friend K. A simulation model of policies directed at tobacco treating tobacco use and dependence. *Medical Decision Making* 2002;22:6-17
15. Levy DT, Ellis JA, Mays D, Huang A-T. Smoking related deaths averted due to three years of policy progress. *Bulletin of the World Health Organization* 2013;91:509-518.
16. Tengs TO, Adams ME, Pliskin JS, et al. Five hundred life-saving interventions and their cost-effectiveness. *Risk Anal* 1995; 15: 369–90.
17. WHO. Policy recommendations for smoking cessation and treatment of tobacco dependence. Tools for public health. Geneva: World Health Organization, 2003.
18. Fiore MC, Jaén CR, Baker TB, et al. Treating Tobacco Use and Dependence: 2008 Update. Clinical Practice Guideline. Rockville, MD: U.S. Department of Health and Human

- Services, Public Health Service.
19. Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ* 2004; 328: 1519.
 20. GBD 2015 Tobacco Collaborators. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990–2015: a systematic analysis from the Global Burden of Disease Study 2015. *Lancet* 2017 April 7;14:48. [http://dx.doi.org/10.1016/S0140-6736\(17\)30819-X](http://dx.doi.org/10.1016/S0140-6736(17)30819-X)
 21. Fidler J, Ferguson SG, Brown J, Stapleton J, West R. How does rate of smoking cessation vary by age, gender and social grade? Findings from a population survey in England. *Addiction* 2013; 108: 1680–85.
 22. International statistical classification of diseases and related health problems, tenth revision (ICD–10). Geneva, World Health Organization, 2007.
 23. US Surgeon General. The Health Consequences of Smoking: Nicotine Addiction: A report of the Surgeon General. Washington DC, Center for Health Promotion and Education, Office on Smoking and Health, 1988.
 24. Tobacco Advisory Group of the Royal College of Physicians. Nicotine addiction in Britain. London, Royal College of Physicians, 2000.
 25. Raw M, Mackay J, Reddy S. Time to take tobacco dependence treatment seriously. *Lancet*, 30 January 2016.
 26. Hughes JR, Keely J, Naud S. Shape of the relapse curve and long term abstinence among untreated smokers. *Addiction* 2004;99: 29-38
 27. West R, Raw M, McNeill A, et al. Health-care interventions to promote and assist tobacco cessation: a review of efficacy, effectiveness and affordability for use in national guideline development. *Addiction* 2015; 110: 1388–403. Available at: http://www.treatobacco.net/en/page_547.php (accessed 7 February 2017)
 28. Bauld L, Hiscock R, Dobbie F, Aveyard P, Coleman T, Leonardi-Bee J, McRobbie H, McEwen A. English stop smoking services: one year outcomes. *International Journal of Environmental Research and Public Health* 2016;13:1175; doi:010,3390/ijerph 13121175
 29. WHO report on the global tobacco epidemic, 2015. Raising taxes on tobacco. Geneva: World Health Organization, 2015.
 30. Borland R, Li L, Driezen P, Wilson N, Hammond D, Thompson ME, Fong GT, Mons U, Willemsen MC, McNeill A, Thrasher JF, Cummings KM. Cessation assistance reported by smokers in 15 countries participating in the International Tobacco Control (ITC) policy evaluation surveys. *Addiction* 2012; 107:197-205
 31. Hatsukami DK, Zeller M, Gupta P, Parascandola M, Asma S. Smokeless tobacco and public health: A global perspective. Washington DC & Atlanta, National Cancer Institute & Centers for Disease Control, 2014
 32. Sinha DN, Agarwal N, Gupta PC. Prevalence of smokeless tobacco use and number of users in 121 countries. *British Journal of Medicine and Medical Research* 2015;9:1-20.
 33. Goodchild M, Perucic AM, Nargis N. Modelling the impact of raising tobacco taxes on public health and finance. *Bulletin of the World Health Organization* 2016;94:250-257.
 34. World Health Organization. Earmarked tobacco taxes, lessons learned from nine countries. Geneva, World Health Organization, 2016
 35. Wilson N, Weerasekera D, Edwards R, Thomson G, Devlin M, Gifford H. Characteristics of smoker support for increasing a dedicated tobacco tax: national survey data from New Zealand. *Nicotine & Tobacco Research* 2010;12:168-73.
 36. World Health Organization. WHO Report on the Global Tobacco Epidemic 2008: the MPOWER package. Geneva, World Health Organization, 2008.
 37. World Health Organization. 19th WHO Model List of Essential Medicines, April 2015. Geneva, World Health Organization.
 38. US Surgeon General. The health consequences of smoking – 50 years of progress: a report

- of the Surgeon General. Atlanta, Department of Health and Human Services, Center for Disease Control and Prevention, 2014.
39. Royal College of Physicians. Nicotine without Smoke. Tobacco harm reduction. London, RCP, 2016.
 40. Chaloupka FJ, Sweanor D, Warner KE. Differential taxes for differential risks – toward reduced harm from nicotine-yielding products. *New England Journal of Medicine* 2015; 373:594-597.
 41. World Health Organization. Strengthening health systems for treating tobacco dependence in primary care. Building capacity for tobacco control: training package. Geneva, World Health Organization, 2013.
http://www.who.int/tobacco/publications/building_capacity/training_package/treatingtobacco_dependence/en/ (accessed 20 Feb 2017)
 42. World Health Organization. Developing and improving national toll-free tobacco quitline services. A WHO manual. Geneva, World Health Organization, 2013.
http://www.who.int/tobacco/publications/smoking_cessation/quit_lines_services/en/ (accessed 20 Feb 2017)
 43. World Health Organization. Toolkit for delivering the 5 As and 5 Rs brief tobacco interventions to TB patients in primary care. Geneva, World Health Organization, 2014.
 44. Tools to implement Article 14: http://www.treatobacco.net/en/page_545.php (accessed 7 February 2017)
 45. Walker N et al. Getting cytisine licensed for use world-wide: a call to action. *Addiction* 2016;111:1895-98.